


IMPLANT SURGERY

Purpose: To replace one or more missing teeth. The Implants are cylinder in form, come in varying lengths and diameters and are either pure titanium or titanium alloy. The Implants replace the missing root or roots. Your dentist will attach an abutment on which a crown will ultimately be placed. This will require an appointment for an impression and a final fitting.

Procedure: This is an in-office procedure. There should be little or no pain during the appointment. The area is anesthetized with local anesthetic and or I.V. sedation. The Implants will be placed under the gums and left for 3-6 months. A second stage procedure will follow to expose the “integrated” healed Implant. ***Once the implant has integrated fully into the jawbone, the next phase involves placement of the implant crown that will be supported by the implant. This is performed by your general dentist or a prosthodontist (dental specialist focused on tooth replacement). The crown or bridge is not included in our fee.***



Pre-Operative Instructions:

1. Eat a normal meal prior to the procedure. If you're having I.V. sedation please refer to the sedation handout for eating instructions.
2. You will be given prescriptions; *normally* an antibiotic, anti-inflammatory and chlorhexidine gluconate rinse. Take as directed on the "Pre-Operative Instructions..." handout.

Post-Operative Instructions

1. Go home, lie down and keep head slightly elevated. Establish that bleeding is controlled, and medications are working. Avoid strenuous activity for the first 48 hours.
2. Expect some swelling this is normal, use an ice pack (provided in post-operative kit) over the surgical area for the remaining portion of the day.
3. After procedure avoid crunchy, hard food for the first 48 hours. When back on a normal diet chew specifically on the other side, if surgery occurred on both sides maintain soft diet for first 2 weeks. Eat well and drink fluids, if you get hypoglycemic or dehydrated you will feel poorly. Even if you do not feel hungry or thirsty, eat and drink anyway. Use of a straw is not recommended for the first 48 hours. This creates negative pressure on the inside of the mouth and can precipitate bleeding. It is advisable not to vigorously rinse spit or swish during this time period. If you wish to rinse put fluid in your mouth and hold in place, turn head from side to side over the sink open your mouth and allow the fluid to drop out. A little pink or redness in the normal volume of saliva is within normal limits.
4. Expect some oozing from the surgical area for the first 48 hours. If any significant bleeding begins (bright red in color greater than normal volume of saliva) which does not stop with gentle pressure, apply a moistened tea bag (not herbal) that has been thinly coated with Vaseline. Put pressure on the bleeding site for approximately 20 minutes. This should stop the bleeding. If this is unsuccessful, call the office for assistance. This means anytime...day or night!
5. The stitched and surgical site will be evaluated in 2 weeks unless otherwise indicated. Do not pull or dislodge any hanging stitches. You may cut them, but do not pull on them.
6. It is very important that nothing is placed over the implant surgical site; i.e., dentures, partials, etc. This area must heal passively for about 3 weeks prior to placing any appliance over the area, unless otherwise specified.
7. Avoid brushing surgical site for at least one week. Gently swab surgical area twice daily with chlorhexidine gluconate rinse (prescription provided).
8. When taking antibiotics and pain medication post operatively stagger the medication, therefore you will not take them at the same time.

NOTE: The quality of the healed result and how long it lasts directly depends on your commitment to daily oral hygiene and continuing care maintenance. It is our goal to never need further intervention.

Remember, you may contact Dr. Wilde at 360-692-0300 or 360-620-2492 at any time if you are having a problem. It is important for you to know that you are not alone and that we are always available to help, reassure or answer any questions.